


DIFCOS Experiment

Doing **I**nteresting, **F**earless, **C**ool and **O**riginal **S**tuff

Fabian Wolf
 QUEST Institute,
 PTB Braunschweig



DIFCOS Team 2024/2025



- Max graduated December 2024 (now at DLR)

DIFCOS Team 2024/2025



New Season



Till Rehmert
(Senior PhD)



Gabriele Gatta
(Guest researcher,
Sias group, Firenze)



2013 Fabian
(dependent group leader)




Sylvain Noël
(Junior PhD)



Johannes Mertens
(Student assistant)


- Max graduated December 2024 (now at DLR)

The dark sector




- Standard model describes only ~ 5 % of the universe

The dark sector




- Standard model describes only ~ 5 % of the universe

Baryon asymmetry




- Standard model CP violation too weak to account for matter anti-matter imbalance

The dark sector




- Standard model describes only $\sim 5\%$ of the universe

Variation of fundamental constants




Baryon asymmetry




- Standard model CP violation too weak to account for matter anti-matter imbalance

The dark sector




- Standard model describes only $\sim 5\%$ of the universe

Variation of fundamental constants




Baryon asymmetry




- Standard model CP violation too weak to account for matter anti-matter imbalance

electron electric dipole moment (eEDM)




The dark sector




- Standard model describes only $\sim 5\%$ of the universe

Variation of fundamental constants




Baryon asymmetry



- Standard model CP violation too weak to account for matter-antimatter imbalance


electron electric dipole moment (eEDM)



Precision spectroscopy of molecular ions

- vibrational spacing scales with m_p/m_e

- internal electric fields of GV/cm possible




Molecule is trapped simultaneously with an atomic ion

- > Laser cooling via the atomic ion
- > Read out the molecule's state via the atomic ion


Motional state serves as transfer bus

Traditional quantum logic spectroscopy




- > Address sideband on optical transition
 - > e.g. Aluminum clock, highly charged ions
- > Dedicated laser for each species
- > Difficult for complex groundstates

Traditional quantum logic spectroscopy




- > Address sideband on optical transition
 - > e.g. Aluminum clock, highly charged ions
- > Dedicated laser for each species
- > Difficult for complex groundstates


Far-detuned Raman quantum logic spectroscopy (FaDeRaQuLoS)






- > pioneered by NIST group for CaH^+
- > Does not require excited metastable state
- > One laser rules them all
- > Required energy gap can be produced by AOMs
 - > Frequency comb allows bridging THz gaps



MgH⁺
















What happened then?

- > New laser had to be set up
- > What to do in the mean time with nicely working setup?

We decided to investigate Ti^+ in the mean time

- > Last year we reported Ti^+ quantum logic spectroscopy


published:

T. Rehmert et al. Quantum logic control of a transition metal ion, Physical Review Letters **134**, 113201 (2025)

g factor measurements using quantum logic and simultaneous co-magnetometry


The grand finale of our unintended excursion into titanium quantum logic spectroscopy

- submitted (T. Rehmert et al. arXiv:2508.15488 (2025))




- Simultaneous Ramsey interrogation
- Sequential readout (Ca: direct, Ti: quantum-logic)

(T. Rehmert et al. arXiv:2508.15488 (2025))



50 Hz magnetic field noise from AC-line

- > **Individual frequencies** affected by magnetic field changes
- > **g factor** not affected
 - > suppression by simultaneous co-magnetometry



(T. Rehmert et al. arXiv:2508.15488 (2025))

In case you forgot: This project is about molecules

Let's get back to molecular ions



*“Beware that, when fighting monsters, you yourself
do not become a monster...
for when you gaze long into the molecule. The
molecule gazes also into you.”*

- > New laser a 1120nm does not dissociate MgH^+

Challenges


	MgH^+	Ti^+
dwelt time in state	seconds	tens of minutes
occupied states @300K	~ 10	4
Zeeman splittings	$\sim \text{kHz}$	$\sim \text{MHz}$

- > New laser a 1120nm does not dissociate MgH^+

Challenges

	MgH^+	Ti^+
dwelt time in state	seconds	tens of minutes
occupied states @300K	~ 10	4
Zeeman splittings	$\sim \text{kHz}$	$\sim \text{MHz}$

Ti^+




- > New laser a 1120nm does not dissociate MgH^+



Challenges



	MgH^+	Ti^+
dwelt time in state	seconds	tens of minutes
occupied states @300K	~ 10	4
Zeeman splittings	$\sim \text{kHz}$	$\sim \text{MHz}$

Ti^+




MgH^+







A new method for measuring heating rates


Not really related to the previous topic




Heating rate measurements










- Phonon arrival time follows **exponential distribution**

- Inverse of mean is heating rate


$$\Gamma_h = \frac{1}{\langle T \rangle} \quad (1)$$

- directly extracted from histogram
- for time-restricted measurements from fit



- > Measurement strength can be tuned by introducing additional deadtime
- > **Incoherent heating** will not be affected
- > **Coherent heating** is suppressed by stronger measurement
 - > Quantum Zeno suppression


SIMULATION NOT MEASURED!



Standard deviation of the mean


- > $\sigma_r = \frac{\Gamma_h}{\sqrt{N}}$
 N : number of detected phonons
- > $\sigma_r = \sqrt{\frac{\Gamma_h}{\tau}}$
 τ : measurement time

Standard quantum limit of heating rate measurements




Ideal Scenario

- > State change only after phonon excitation
- > State change after each phonon excitation




Ideal Scenario

- > State change only after phonon excitation
- > State change after each phonon excitation




Spontaneous decay

- > Without heating ion goes into bright state
- > Overestimation of heating rates




Ideal Scenario

- > State change only after phonon excitation
- > State change after each phonon excitation




Spontaneous decay

- > Without heating ion goes into bright state
- > Overestimation of heating rates



Sideband infidelities

- > Despite heating population remains in dark state
- > Underestimation of heating rates




Auto-calibrating

- > No initial guess of heating rate requires
- > Measurement time is automatically adapted

Broad range


- > Covers typical heating rates
- > Depends on ion species

Metrologically clean



Todo list

- > Measure Quantum Zeno suppression for coherent heating
- > Investigate correlation of heating with 50Hz noise
 - > similar to photon-correlation for micro-motion measurements
- > Model systematic uncertainties



Ground loops?

